

APPENDIX 2

EARDLEY ESCARPMENT
DESCRIPTION AND CONSERVATION ISSUES

DEFINITION OF AN ESCARPMENT

In geology, an escarpment is a transitional zone between different physio-geographic provinces that involves a steep elevation differential characterized by a cliff. More commonly, an escarpment is a transition from one series of sedimentary rocks to another series of a different age and composition. It is usually the result of loss of the newer rock (situated on top of the older rock) through erosion.

Some escarpments were formed by tectonic movement (deforming movements of the Earth's crust or volcanic activity), and in particular by the vertical movement of the Earth's crust along a fault, creating a fault scarp. A fault is a fracture in the ground in which the adjacent surfaces are displaced along the plane of the fracture.

An escarpment can be divided into several portions, each subject to different levels of natural and human-induced impacts.

The first portion is the summit or roof (the upper surface of the escarpment, or the land immediately above it), and the foot or base (the land located at the bottom, covered with piles of rocks that have fallen from the cliff during erosion). It is here, at the summit and base, that the most abundant vegetation is found, in a variety of layers.

Between the two, the vertical or sloping portion of the escarpment is known as the wall. This is the portion that is most exposed to natural elements such as wind and rain, and usually has a large number of cracks, crevasses and ledges. A ledge or platform is a natural protrusion above a steep slope, formed by a resistant layer, around a plateau, with a softer layer below. Often, a thin layer of soil builds up on the ledge, allowing for the development of plant life. Cracks are discontinuities, breaks or tabular openings in the rock, ranging from a few centimetres to several decimetres in length, and from a few millimetres to several centimetres in width. They are produced by physical constraints. Crevasses, for their part, are superficial openings in the rock, sometimes used by plant or animal species as shelter.

ESCARPMENT RARITY, FRAGILITY AND WEALTH

Eardley Escarpment is located in the Province of Québec, and runs along the line between the Canadian Shield and the St. Lawrence Lowlands. It is also the richest and most fragile environment in Gatineau Park.

Eardley Escarpment is a cliff lying along a south-south-west line. It is approximately 300 metres high, with an average height of more than 200 metres, and is the dominant topographical element in the Outaouais region. It begins in the City of Gatineau and runs north-eastwards along the Ottawa River for several dozen kilometres, forming a characteristic rock slope alignment.

This type of environment is extremely rare; there are only a handful of documented escarpments in Canada and the world.

According to Gagnon (1981), Eardley Escarpment's ecological value for conservation is very high due to the diversity of its plant communities. In addition, its red oak (*Quercus rubra*), white oak (*Quercus alba*) and red oak/red cedar (*Juniperus virginiana*) stands are some of the rarest vegetation communities in the Ottawa Valley and Québec.

The plant species at risk are mainly associated with the Escarpment's carbonate rock outcrops and open oak stands, where their principal concentrations are found (Lavoie, 1992). These species are often vulnerable; nearly 40% exhibit limited distribution, require specialized habitats and have small populations.

Because of the Escarpment's micro-climate, it has developed a form of plant life that is unusual in Western Québec. The conditions on the Escarpment slopes are closer to those found further south, for example in the American Mid-West. The warm micro-climate is conducive to the growth of certain specific species such as the blunt-lobe cliff fern (*Woodsia obtusa*), the walking fern (*Asplenium rhizophyllum*), the white oak, the red cedar and the common hackberry (*Celtis occidentalis*). In many places, the vegetation resembles a savannah, with tall grasses and sparse trees. The slow growth and underdeveloped aspect of the trees, some of which are more than 100 years old, are evidence of the difficult edaphic conditions on the Escarpment.

Other threatened and vulnerable species, as well as species likely to be designated as such, are found in the Eardley Escarpment ecosystem. Gagnon et al. (1993) studied the population dynamics of eight threatened or vulnerable plant species, including the woodland sunflower (*Helianthus divaricatus*), Douglas' knotweed (*Polygonum douglasii*) and the fragrant sumac (*Rhus aromatica*), which are vulnerable species in Québec. Gagnon and Hay (1986) studied the

eastern few-fruit sedge (*Carex oligocarpa*), this being only the seventh known location in Canada of this species at risk.

The Escarpment, which is the only place within Gatineau Park where the red cedar is found, contains the largest population of red cedar in Québec, estimated at 15,000 trees (NCC, 2002c) and accounting for 80% of the total population in Québec (Forest, 1994). According to Forest (1994), the Eastern red cedar is one of the rarest species of trees in Québec, found almost exclusively within the oak-red cedar stands on Eardley Escarpment. This is a very rare type of community within the province, found only in the Ottawa Valley. Because the red cedar populations are located mainly on the Escarpment walls, they are under direct pressure from rock climbing activities. The most popular rock climbing site is also home to the second largest concentration of red cedar in the Park.

The other benefits of the Escarpment's micro-climate include a number of bird species above the cliffs. The birds take advantage of rising air currents and warm winds directed upwards by the Escarpment's topography. Birds of prey migrate through the region in spring and early fall. During these periods, it is possible to observe the red-tailed hawk (*Buteo jamaicensis*) and the turkey vulture (*Cathartes aura*). Many species use the cliff for nesting, and groups of crows and rooks are regularly observed by climbers. Owls, rarely seen because of their nocturnal habits, are also common on the Escarpment. There are several potential nesting sites for the American peregrine falcon (*Falco peregrinus anatum*) on the Eardley Escarpment. These nesting sites are monitored every five years by the MDDEP.

Land-based wildlife is also abundant on the Escarpment. For example, the cervids winter on the western portion of the Escarpment. During March and early April, it is not unusual to see dozens of animals feeding in the fields at the base of the Escarpment, or travelling the network of paths they create by trampling the snow covered slopes. In winter, packs of wolves (*Canis lupus lycaon*) visit the Escarpment, hunting for white-tailed deer (*Odocoileus virginianus*). This part of the Park is also inhabited by black bear (*Ursus americanus*), which can be observed on occasion. The bears are attracted by the plentiful supplies of berries growing on top of the hills. Their dens are usually located alongside the Escarpment crest. Other common species include the racoon (*Procyon lotor*), muskrat (*Ondatra zibethicus*), squirrel (*Sciurus carolinensis*, *Tamiasciurus hudsonicus*) and chipmunk (*Tamias striatus*).

In 1990, a juniper hairstreak (*Callophrys gryneus*), a butterfly often found around the red cedar, was captured for the first time in Québec at the bottom of Eardley Escarpment. It is now classified as a threatened species in Québec by the Société d'entomologie du Québec (SEQ). Insect populations associated with the red cedar and other Escarpment plants have been studied on numerous occasions, highlighting the entomological richness of this habitat (Landry, 1990). In 1991, new lepidoptera species were seen along the Escarpment (Landry and Landry, 1991), while in 2001, researchers found six species of beetles that had never before been documented in Québec (Laplante, 2001). Goulet (1994) also found the first examples in Québec of three hymenoptera families, linked to the presence of the red cedar. The discovery of these species in Québec highlights the unique character of this region's wildlife.

Because of all these factors, Eardley Escarpment is without question Gatineau Park's richest natural environment. Because of its hot, dry climate and steep slopes, it is also the most fragile, and is particularly sensitive to erosion.

IMPACTS OF ROCK CLIMBING ON THE HOST ENVIRONMENT, AND CONSERVATION OPTIONS

A study by Dubé (1995) showed the significant impact of climbing on some Eardley Escarpment habitats. Many other studies have also been carried out, some on the question of protecting plant species and others on damage to the rock face and the potential decline in the number of wild animal species. In all, six major impacts have been identified. The table below summarizes the available information. In addition, Kuntz and Larson (2005) report that the impact of rock climbing varies according to the degree of difficulty of the wall. The study recommends the use of walls with a level of difficulty higher than 5.10, rather than easier walls, since their impact on the environment is less severe.

The conservation options set out in the table have been applied in some locations, and it is therefore possible to comment on them:

Closure of climbing sites: This management option has frequently been proposed in the literature. It allows plant and animal species to re-colonize the environment, increases species wealth, protects the plant cover and avoids loss of essential habitats. However, the literature also shows that it often does not produce satisfactory results because climbers continue to use the prohibited sites, and the closures are difficult to enforce. Even so, the results observed on Mont Saint-Hilaire are encouraging. (Camp and Knight, 1998; Rusterholz et al., 2004; Kelly and Larson, 1997; McMillan and Larson, 2002)

Closure of certain climbing paths and installation of permanent paths: This option, too, is mentioned frequently in the literature. It allows users to continue their activity, except during critical periods of the year, for example during flowering or nesting. The literature shows that climbers tend to comply with this type of rule. A management plan can also be introduced, allowing those in charge to set limits on climbing and introduce measures to ensure environmental sustainability (Camp and Knight, 1998; Rusterholz et al., 2004)

Introduction of monitoring programs: Also cited frequently in the literature, this option identifies damage to the ecosystem and allows for ongoing observation of the impacts of rock climbing and recreational activities on the cliffs and related communities. Based on these observations, managers are able to assess the intensity of the activities and decide on a management approach. A monitoring program may be combined with complete or partial closure of climbing sites. (Rusterholz et al., 2004; Richardson, 1999; McMillan and Larson, 2002)

Status quo: Many of the reports cited in this document show that the environment is affected significantly by rock climbing activities on the Escarpment cliff, and that measures are needed to correct this situation. In addition, it has been shown that the environment's level of resilience declines steadily as a result of species loss and community impoverishment, preventing the ecosystems from resuming their functions and developing normally after disturbance.

IMPACTS ON THE ENVIRONMENT	CONSERVATION OPTIONS
REDUCTION IN SPECIES WEALTH	
<p>Decline in plant and animal species (Baker, 1999; Rusterholz et al., 2004; CPAWS, 2005).</p> <p>Decline in lichen species on climbing sites, compared to non-climbing sites (Farris, 1998).</p> <p>Decline in tree, bush and non-graminaceous grass species on climbing sites, compared to non-climbing sites (Camp and Knight, 1998).</p> <p>Decline in snail population wealth (McMillan et al., 2003).</p> <p>More bird species (60 %) on non-climbing sites (Krajick, 1999).</p> <p>Loss of diversity in bryophyte, moss and liverwort species (McMillan and Larson, 2002; Kingsley, 2002).</p> <p>Health considered severe for three of the seven red cedar species (NCC, 2002a).</p> <p>Decline in the number of three calcicole fern species: the walking fern (<i>Asplenium rhizophyllum</i>), purple cliffbreak (<i>Pellea atropurpurea</i>) and bluntlobe cliff fern (<i>Woodsia obtusa</i>) (Gagnon, 2002).</p> <p>Significant differences in the number of species and relative frequency of many species (Farris, 1995).</p> <p>Substantial damage to vegetation in areas used by rock climbers (Genetti and Zenone, 1987).</p>	<p>Close or control of access paths and inform users (Camp and Knight, 1998; Cornish, 2004; Farris, 1998; Francis, 2001; Kelly and Larson, 1997; McMillan and Larson, 2002; Nuzzo, 1995; Richardson, 1999; Rusterholz et al., 2004)</p>
A DECLINE IN PLANT COVER	
<p>Decline of more than 5% on climbing sites, compared to non-climbing sites (Rusterholz et al., 2004).</p> <p>Greater density of large white cedars on non-climbing sites (Farris, 1998).</p> <p>Decline in lichen cover (Krajick, 1999).</p> <p>Decline in plant cover at the base of climbing sites (McCarthy, 2003).</p> <p>Less abundant plant life (Baker, 1999).</p> <p>Significant differences in total vegetation cover (Farris, 1995).</p> <p>Vegetation shift and loss of plant growth sites (Genetti and Zenone, 1987).</p>	<p>Reduce access during critical (breeding) periods and install permanent path systems (Camp and Knight, 1998; Farris, 1998; Francis, 2001; Kelly and Larson, 1997; McMillan and Larson, 2002; Nuzzo, 1995; Rusterholz et al., 2004)</p>
SOIL EROSION	
<p>Soil compaction (McMillan et al., 2003).</p> <p>Soil displaced by climbers (Francis, 2001).</p> <p>Increase in soil degradation (Cornish, 2004).</p> <p>Creation of deep run-off runnels in the soil (Genetti and Zenone, 1987).</p> <p>Signs of soil erosion (CPAWS, 2005).</p>	<p>Limit access to climbing sites and prohibit the use of pitons and carabiners (Camp and Knight, 1998; Farris, 1998; Francis, 2001; Rusterholz et al., 2004; Cornish, 2004)</p>
EROSION OF CLIFF WALLS	
<p>Physical signs of damage on cliff walls (Francis, 2001; Cornish, 2004; Kelly and Larson, 2004).</p> <p>Evidence of rain-resistant chalk on the rock walls (McCarthy, 2003).</p> <p>Signs of cliff wall erosion (CPAWS, 2005).</p>	<p>Prohibit the use of pitons and carabiners, reduce treading and inform users (Camp and Knight, 1998; Farris, 1998; Kelly and Larson, 1997; Rusterholz et al., 2004; Cornish, 2004)</p>

IMPACTS ON THE ENVIRONMENT	CONSERVATION OPTIONS
ALTERATION OF PLANT AND ANIMAL COMMUNITY COMPOSITION	
<p>Change in the frequency of dwarf trees such as the creeping germander (<i>Teucrium chamaedrys</i>) and mountain germander (<i>Teucrium montanum</i>) (Rusterholz et al., 2004).</p> <p>Increase in monocotyledon populations on sites used by climbers (Farris, 1998).</p> <p>Few trees in younger age classes (Kelly and Larson, 1997).</p> <p>Change in snail population composition (McMillan et al., 2003).</p> <p>Invasions of invasive species on sites used by climbers (Oosthoek, 2002).</p> <p>Different bird communities, altering spatial distribution (Camp and Knight, 1998).</p> <p>Changes in the number of invasive bird species (Camp and Knight, 1998).</p>	<p>Introduce monitoring programs (Camp and Knight, 1998; Farris, 1998; Francis, 2001; Kelly and Larson, 1997; McMillan and Larson, 2002; Nuzzo, 1995; Richardson, 1999; Rusterholz et al., 2004)</p>
HABITAT LOSS	
<p>Large percentage of bare rock (Rusterholz et al., 2004).</p> <p>Decline in the number of growth sites suitable for plants (Camp and Knight, 1998).</p> <p>Degradation of the red cedar (<i>Juniperus virginiana</i>) habitat (NCC, 2002a).</p>	<p>Close or control access and inform users (Camp and Knight, 1998; Cornish, 2004; Farris, 1998; Francis, 2001; Kelly and Larson, 1997; Nuzzo, 1995; Richardson, 1999; Rusterholz et al., 2004)</p>

BIBLIOGRAPHY

- ACC : ALPINE CLUB OF CANADA, 2004. *A submission by the Alpine Club of Canada to the National Capital Commission in response to the Preliminary Gatineau Park Master Plan*, Protecting Gatineau Park: the need for balance and cooperation.
- ACCESS FUND. *Raptors and climbers : how climbing restrictions are set up and managed*, Climbing and natural resources management.
- ACCESS FUND, 1999. *Restoration, conservation and policy proposal for rock climbing – McConnells Mill State Park, Pennsylvania*, Climbing and natural resources management.
- ACCESS FUND, 2000. *Climbing management: a guide to climbing issues and the production of a climbing management plan*, Climbing and natural resources management.
- ATTARIAN, A., 1992. *An investigation of the ecological and social impacts caused by rock climbers*, In C. Rademacher & R. Watters (Eds.), Proceedings of the 1991 International Conference on outdoor recreation and education, Idaho State University Press, 7-16.
- ATTARIAN, A. ET K. PYKE, 2000. *Climbing and natural resources management : an annotated bibliography*, North Carolina State University and the Access Fund.
- BAKER, B., 1999. *Controversy over use of rock-climbing anchors may be missing the mark*, BioScience, 49, 529.
- BARNARD, C., 1986. *Application of the recreation opportunity spectrum to Gatineau Park*, National Capital Commission for Gatineau Park, Visitor Services & Planning & Development Sections.
- BLACKWELL, S.D., 1999. *Environmental impacts of rock climbing in Yosemite Valley, Yosemite National Park*, San Diego State University.
- BOISE CLIMBER'S ALLIANCE ET IDAHO DEPARTMENT OF FISH AND GAME, 1999. *Guidelines for protecting cliff-nesting raptors and climbing management at Black Cliffs, Boise, Idaho*, Idaho Department of Fish and Game, Nampa.
- CAMP, R.J. ET R.L. KNIGHT, 1998. *Effects of rock climbing on cliff plant communities at Joshua Tree National Park, California*, Conservation Biology, Volume 12 Issue 6 Page 1302 Décembre 1998.
- CAMP, R.J. ET R.L. KNIGHT, 1998. *Rock climbing and cliff bird communities at Joshua Tree National Park, California*, Wildlife Society Bulletin, 26(4): 892-898.
- CENTRE D'INTERPRÉTATION DE LA GÉOLOGIE DE GRENVILLE, 2003. Sites et circuits du patrimoine naturel de la région de l'Outaouais, p 30-34.
- COMMISSION DE LA CAPITALE NATIONALE, 2003. *Dossier escalade: Sommaire des résultats des études environnementales*, Parc de la Gatineau, Parois rocheuses de l'escarpement d'Eardley.
- COMMISSION DE LA CAPITALE NATIONALE, 2007. *Escarpement d'Eardley*. [En ligne] http://www.canadascapital.gc.ca/bins/ncc_web_content_page.asp?cid=16297-16299-10170-16394-16384&lang=2&bhcp=1.
- CORNISH, M., 2004. *A strategy of accommodation for Devils Tower National Monument*, Washington DC.
- COMITÉ SUR LA SITUATION DES ESPÈCES EN PÉRIL AU CANADA (COSEPAC), 2000. *Mise à jour – Évaluation et Rapport de situation du COSEPAC sur le ginseng à cinq folioles (Panax quinquefolius) au Canada*, Espèce en voie de disparition 2000.
- DE BENEDETTI, S., 1990. *Impacts of rock climbing and mitigation actions taken at Pinnacles National Monument*, paper presented at the George Wright Society Conference on research and resource management in parks and public lands.
- DRIESE, K.L. ET D.A. ROTH, 1992. *A description of the vascular flora and mammal fauna and the effects of human disturbance on the summit of Devils Tower*, University of Wyoming, Laramie.

- DUBÉ, 1994-1995. *Activité d'escalade à l'intérieur du Parc de la Gatineau*. Étude réalisée par la Commission de la Capitale Nationale.
- DUPUIS V., 2001. *Suivi environnemental de sentiers autorisés et non autorisés pour la pratique du vélo de montagne*, Préparé pour la Commission de la Capitale Nationale, Parc de la Gatineau, Gestion des ressources naturelles et des terrains.
- FARRIS, M.A., 1995. *The effects of rock climbing on the cliff flora of three Minnesota State Parks*, Conservation Biology Research Grants Program, Division of Ecological Services, Final report to the Minnesota department of natural resources.
- FARRIS, M. A., 1998. *The effects of rock climbing on the vegetation of three Minnesota cliff systems*, Department of Biology, Hamline University, USA, *Canadian Journal of Botany*, 76: 1981-1990 (1998).
- FINDLAY, S., J. MCDONALD, S. IRVEN ET J. HALL, 2005. *Gatineau Park Conservation Management Plan – Rock Climbing*, Department of Biology, University of Ottawa.
- FINDLAY, S., T. STANLEY ET A. WASIAK, 2005. *Gatineau Park Conservation Management Plan – Operational Issues and human impacts of hiking*, Department of Biology, University of Ottawa.
- FOREST, G., 1994. *Dynamique des populations de genévrier de Virginie de l'escarpement d'Eardley, parc de la Gatineau*, Université du Québec à Montréal.
- FRANCIS, W., 2001. *Rock climbing is damaging cliff-dwelling plants in Red River Gorge*, *Kentucky Native Plant Society*, 16, 1-3.
- GAGNON, D., 1978. *Ecological assessment of the Eardley Escarpement*, Université du Québec à Montréal.
- GAGNON, D. ET A. BOUCHARD, 1981. *La végétation de l'escarpement d'Eardley, parc de la Gatineau, Québec*, *Canadian Journal of Botany*, 59 (12): 2667-2691 (1981).
- GAGNON, D. ET G. HAY, 1986. *Carex oligocarpa (Cyperaceae), a rare sedge in Canada newly discovered in Québec*, Université du Québec à Montréal.
- GAGNON, D., L. COLLINS, P. NANTEL, G. FOREST ET N. LAVOIE, 1993. *Étude de la dynamique des populations de huit espèces de plantes menacées ou vulnérables du Québec*, Université du Québec à Montréal.
- GAGNON, D., 2002. *Caractéristiques du micro habitat de trois espèces de fougères calcicoles du parc de la Gatineau susceptibles d'être désignées menacées ou vulnérables*, Université du Québec à Montréal.
- GENETTI, C. ET P. ZENONE, 1987. *The effect of rock climbers on the environment at Pinnacles National Monument, Monterey and San Benito counties, California*, Technical report No. 27, Cooperative National Park Resources Studies Unit, University of California at Davis.
- GOULET, H., 1994. *Première mention de trois familles d'hyménoptères au Québec : AMPULICIDAE, EVANIIDAE et ROPRONIIDAE*, Ministère Agriculture Canada.
- GRAHAM, L. ET R.L. KNIGHT, 1999. *Bird and plant communities associated with cliffs in Jefferson County, Colorado*, Colorado State University, Department of fishery and wildlife biology, Fort Collins.
- JODICE, P. ET K. PYKE, 1999. *Climbing and cliff ecology*, *Science*, 284:5413, 433.
- KELLY, P.E. ET D.W. LARSON, 1997. *Effects of rock climbing on populations of pre-settlement eastern white cedar on cliffs of the Niagara*, *Conservation Biology*, 11 (5), 1125-1132.
- KELLY, P.E., 1998. *Environmental stewardship along the Niagara Escarpment*, *The Canadian Alpine Journal*, 81, 79-81.
- KINGSLEY, D., 2002. *Scaling the damage of rock climbing*, *News in Science*, April 2002.
- KNOLL, M., 2001. *Soil compaction due to rock climbing and hiking activity at cliff margins on the Southern Cumberland Plateau, Tennessee*, University of the South, Department of forestry and geology, Sewanee.
- KRAJICK, K., 1999. *Scientists and climbers discover cliff ecosystems*, *Science*, 283, 1623-1625.

- KUNTZ, K.L. et D.W. LARSON, 2005. *Influences of Microhabitat Constraints and Rock-Climbing Disturbance on Cliff-Face Vegetation Communities*. Conservation Biology, Volume 20, No. 3, 821-832.
- LANDRY, -F. ET B. LANDRY, 1991. *Mentions nouvelles ou intéressantes de Lépidoptères dans le sud du Québec en 1991*, Ministère Agriculture Canada.
- LANDRY, B., 1990. *Recherche entomologique, saison 1990*, Université de Carleton.
- LAPLANTE, S., 2001. *Capture en Outaouais de quelques coléoptères rarement trouvés ou nouveaux pour la faune du Québec*, Ministère Agriculture Canada.
- LARSON, D.W., U. MATTHES ET P. KELLY, 1999. *Cliff as natural refuges*, American Scientist, 411-417.
- LAVOIE, G., 1992. *Plantes vasculaires susceptibles d'être désignées menacées ou vulnérables au Québec*, Ministère de l'Environnement du Québec.
- MALKIN, D.R., 2000. *Effects of rock climbing on populations of Silene seelyi, a rare perennial plant*, Master's thesis, University of Washington, Seattle.
- MALOUIN, 2002a. *Évaluation de l'état de sept populations de genévriers de Virginie localisées sur l'escarpement d'Eardley*. Étude réalisée par la Commission de la capitale nationale.
- MALOUIN, 2002b. *Évaluation des répercussions environnementales liées à la pratique de l'escalade sur cinq parois rocheuses situées sur l'escarpement d'Eardley*. Étude réalisée par la Commission de la Capitale Nationale.
- McCARTHY, L.P., 1993. *National Parks grapple with rock climbing*, National Parks, 67, 22.
- McCARTHY, T., 2003. *Wearing down the mountains: are rock climbers leaving cliff faces in less than peak conditions?*, Time, 162, 54.
- McMILLAN, M.A. ET D.W. LARSON, 2002. *Effects of rock climbing on the vegetation of the Niagara escarpment in southern Ontario, Canada*, Conservation Biology, 16: 389-398, 2002.
- McMILLAN, M.A., J.C. NEKOLA ET D.W. LARSON, 2003. *Effects of rock climbing on the land snail community of the Niagara Escarpment in Southern Ontario, Canada*, Conservation Biology, 17: 616-621.
- MORGAN, K., 2003. *At a snail's place: rock climbing cuts mollusc diversity*, Science News Online, Vol. 163, No. 15, april 12, 2003.
- NUZZO, V.A., 1995. *Effects of rock climbing on cliff goldenrod (Solidago sciaphila Steele) in northwest Illinois*, American Midland Naturalist, 133: 229-241.
- NUZZO, V.A., 1996. *Structure of cliff vegetation on exposed cliffs and the effect of rock climbing*, Canadian Journal of Botany, 74: 607-617.
- OVERLIN, A., G.W. CHONG, T.J. STOHLGREN ET J. RODGERS, 1999. *Vertical Veg: Partnership project to address resources protection and quality visitor experience in rock climbing areas of Joshua Tree National Park*, United States Department of the Interior, National Park Service, Joshua Tree National Park, Twentynine Palms, in cooperation with the Access Fund and the California Native Plant Society.
- PARC DE LA GATINEAU, 2004. *Synthèse écologique*, textes synthèses.
- RÉSERVE NATURELLE HAUTS DE CHARTREUSE, 2006. *Gestion de la pratique de l'escalade sur la Réserve Naturelle des Hauts de Chartreuse*, Groupe de travail. [En ligne] http://www.promo-grimpe.com/doc_pdf/convention_pnrc.pdf
- RICHARDSON, H., 1999. *Threats posed by rock climbers to birds nesting on cliffs in the South Okanagan*, In proceedings of a Conference on the Biology and Management of Species and Habitats at Risks, Kamloops, B.C., 1,490.
- ROCKY MOUNTAIN NATIONAL PARK, 1990. *Task force findings : climbing in Rocky Mountain National Park*, Rocky Mountain National Park, Estes Park.
- RUSTERHOLZ, H.P., S.W. MULLE ET B. BAUR, 2004. *Effects of rock climbing on plant communities on exposed limestone cliffs in the Swiss Jura mountains*, Applied Vegetation Science, 7, 35-40.

- SCHUSTER, R.M., J.G. THOMPSON ET W.E. HAMMIT, 2001. *Rock climbers' attitudes toward management of climbing and the use of bolts*, Environmental Management, 28:3, 403-412.
- SPEAR, P.W. ET M.J. SCHIFFMAN, 1979. *Rock climbing and endangered plants : a case study*, In proceedings of the speciality conference, National conference on recreation planning and development, pp. 630-636.
- SPRUNG, G., 2004. *A summary of scientific studies that compare mountain biking to other forms of trail travel*, International Mountain Bicycling Association.
- SYNDICAT DE COOPÉRATION POUR LE PARC NATUREL RÉGIONAL DES VOSGES DU NORD, 1997. *Charte pour la pratique de l'escalade sur les rochers des Vosges du Nord*, Charte signée le 6 décembre 1997 par la Fédération Française de la Montagne et de l'Escalade, l'Association S.O.S. Faucon pèlerin, l'Office National des Forêts, les groupements forestiers concernés et le Sycoparc.
- TROUTET, Y., 1977. *Climbing on the Eardley Escarpment*.
- WILD, M. ET D. GAGNON, 2001. *Prédire l'habitat d'une fougère rare au parc de la Gatineau, Québec, à l'aide d'un modèle GIS basé sur les données de présence*, Université du Québec à Montréal.
- ZSCHOKKE, S., C. DOLT, H-P. RUSTERHOLZ, P. OGGIER, B. BRASCHLER, G.H. THOMMEN, E. LUDIN, A. ERHARDT ET B. BAUR, 2000. Short-term responses of plants and invertebrates to experimental small-scale grassland fragmentation, *Oecologia* (2000) 125: 559-572.

WEBOBGRAPHY

- <http://utime.unblog.fr/2007/03/30/la-biodiversite-lautre-choc/>
- http://fr.wikipedia.org/wiki/R%C3%A9silience_%C3%A9cologique
- http://www.ecologie.gouv.fr/IMG/pdf/CSPNB_BiodivExemples_38-61.pdf
- <http://www.canadianencyclopedia.ca/index.cfm?PgNm=TCE&Params=f1ARTf0002641>
- <http://en.wikipedia.org/wiki/Escarpment>
- http://en.wikipedia.org/wiki/Niagara_Esacarpment
- [http://en.wikipedia.org/wiki/Onondaga_\(geological_formation\)](http://en.wikipedia.org/wiki/Onondaga_(geological_formation))
- http://en.wikipedia.org/wiki/Devil's_Rock
- http://geoscape.nrcan.gc.ca/toronto/pdf/escarp_activ3_f.pdf
- <http://fototek.geol.u-psud.fr/~v.ansan/ObjectifTerre/Geomorphologie/ Escarpement/escarp.htm>
- http://www.canadacapital.gc.ca/bins/ncc_web_content_page.asp?cid=16297-16299-10170-49685-49721-59486&lang=2&bhcp=1
- <http://www.acfas.ca/CONGRES/congres65/S2143.htm>
- <http://www.centrenature.qc.ca/pdf/Faucon/Faucon0905.pdf>
- <http://www.centrenature.qc.ca/conservation/montagne.html#objectifs>